



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

SUPPLEMENTAL LABEL

*This supplemental label **expires on 03/02/2019** and must not be used or distributed after this date.*

ILeVO®

FOR USE ON: Soybean

ACTIVE INGREDIENT:

FLUOPYRAM: N-[2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl]-2-(trifluoromethyl)benzamide*..... 48.4%

OTHER INGREDIENTS:..... 51.6%

TOTAL: 100.0%

Contains 5 lbs FLUOPYRAM per gallon (600 g FLUOPYRAM per liter)

*(CAS Number 658066-35-4)

EPA Reg. No. 264-1167

EPA Est. No.

SUSPENSION CONCENTRATE

KEEP OUT OF REACH OF CHILDREN CAUTION

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of FLUOPYRAM. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect seed spilled during loading

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This Supplemental Label and the entire ILeVO container label, EPA Reg. No 264-1167, must be in possession of the user at the time of application.
- Read the label affixed to the container for ILeVO before applying.
- Use of ILeVO according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for ILeVO. Refer to the ILeVO container label for additional instructions and restrictions.

ROTATIONAL CROP RESTRICTIONS

The following crops can be replanted immediately following the last application of Fluopyram:
Artichoke, globe; Berry, low growing, except cranberry, subgroup 13-07G; Brassica, Head and Stem, subgroup 5A; Brassica Leafy Greens, subgroup 5B; Bushberry subgroup 13-07B; Caneberry subgroup 13-07A; Carrot; Cherry subgroup 12-12A; Corn, field, grain; Corn, pop, grain; Corn, sweet kernel plus cob with husks removed; Cottonseed subgroup 20C; Dill seed; Fruit, citrus, group 10-10; Fruit, pome, group 11-10; Fruit, small vine climbing, except fuzzy kiwifruit, subgroup 13-07F; Ginseng; Grain, cereal, group 15 (except corn) which include: barley, buckwheat, millet (pearl and proso), oats, rice; rye, sorghum, teosinte, triticale, and wheat; Herb subgroup 19A; Hop, dried cones; Leafy greens subgroup 4A; Leaf petioles subgroup 4B (except watercress); Legume Vegetables (except cowpea and dried peas); Melon subgroup 9A; Nut, tree, group 14-12; Onion, bulb, subgroup 3-07A; Onion, green, subgroup 3-07B; Peach subgroup 12-12B; Plum subgroup 12-12C; Peanut; Pepper/eggplant subgroup 8-10B; Rapeseed subgroup 20A; Soybean; Squash/Cucumber subgroup 9B; Sugarbeet, roots; Sunflower subgroup 20B; Tomato subgroup 8-10A; Vegetable, root, except sugarbeet, subgroup 1B; Vegetable, tuberous and corm, subgroup 1C.

Alfalfa can be planted after 14 days and sugarcane can be planted after 14 days in Region 3. Do not rotate to crops other than those listed above.

Do not rotate to crops other than those listed above.

SOYBEAN		
Disease Control	Application Rates	Product Instructions
Sudden Death Syndrome (<i>Fusarium viguliforme</i>)	0.15 - 0.25 mg ai/seed or 1.18 to 1.97 fl oz (35 to 58 ml) / 140,000 soybean seeds	Apply using commercial slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Dilute product with sufficient water to ensure complete seed coverage. Apply to high quality, properly cleaned seeds.
Soilborne Nematodes (<i>Heterodera glycines</i> , <i>Meloidogyne incognita</i> , <i>Rotylenchulus reniformis</i> , <i>Pratylenchus</i> spp., <i>Hoplolaimus</i> sp.)	0.075 - 0.25 mg ai/seed or 0.6 to 1.97 fl oz (18 to 58 ml)/ 140,000 soybean seeds	This product does not contain colorant. The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, part 2.25. Any colorant added to treated seed must be cleared for use under 40CFR, part 153.155.
Restrictions:		
<ul style="list-style-type: none"> • Seeds may not be broadcast planted. • Regardless of formulation or method of application soil, foliar or seed treatment, do not apply more than 0.446 lbs of fluopyram per acre per year. • Do not allow livestock to graze soybean forage or hay for 7 days after application and do not harvest soybean forage or bean hay for food or feed for 7 days after application. 		

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

SAL on 03/02/2016

ILeVO is a Registered trademark of Bayer.